

ON THE (NON) EXISTENCE OF VISCOSITY SOLUTIONS OF MULTI-TIME HAMILTON-JACOBI EQUATIONS

A. DAVINI, M. ZAVIDOVIQUE

ABSTRACT. We prove that the multi-time Hamilton-Jacobi equation cannot be solved in the viscosity sense, in the non-convex setting, even when the Hamiltonians are in involution.

INTRODUCTION

There are two important Hamilton-Jacobi equations associated with a continuous function H defined on the cotangent bundle of a smooth Riemannian manifold M : the stationary equation, that is a nonlinear PDE of the kind

$$H(x, d_x u) = c \quad \text{in } M,$$

where c is a real constant and the unknown u is a real function defined on M ; and the evolutionary equation, which is a time-dependent equation of the form

$$\begin{cases} \partial_t u + H(x, d_x u) = 0 & \text{in } (0, +\infty) \times M \\ u(0, x) = u_0(x), & \text{in } M, \end{cases} \quad (1)$$

where $u_0 : M \rightarrow \mathbb{R}$ is an initial datum, and the unknown u is a real function of variables $(t, x) \in [0, +\infty) \times M$.

The first problem that naturally arises is that of finding a good notion of solution. It is in fact well known that such equations do not usually admit classical solutions. For instance, the method of characteristics provides, for smooth initial data and under general assumptions on H , classical solutions for (1) only for small times, until shocks between characteristics occur.

A first idea is to look for Lipschitz functions which solve the equation almost everywhere. However, such a notion of solution is inadequate since it lacks of good uniqueness and stability properties, see for instance [Bar94]. That is why two better notions of weak solutions were introduced.

The first one, introduced by Crandall and Lions in 1983 [CL83], is that of *viscosity solution* and it applies to a wide range of first and second order nonlinear PDE. It is in some way reminiscent of distributions, in the sense that it uses test functions to drop derivatives, and permits to name solutions functions that are just continuous, regardless to their regularity. This notion has revealed to be extremely powerful and flexible, and it has been generalized since then in many different directions.

The second one, more geometric in nature, is termed *variational solution* and was introduced by Chaperon, Sikorav and Viterbo (see [Cha91] or [CV08] and references therein) and is specific of evolutionary equations of the kind of (1). It follows the

2010 *Mathematics Subject Classification.* 35F21, 49L25, 37J50.

Key words and phrases. commuting Hamiltonians, viscosity solutions, symplectic geometry.

The second author is financed by ANR-12-BLAN-WKBHJ.

idea of the method of characteristics, taking the graph of the differential of the initial condition and pushing it with the Hamilton flow. When the image is no longer a graph, a natural way to cut the obtained Lagrangian manifold, known as the graph selector, allows to reconstruct a function which is the desired variational solution.

Note that, as can be expected, both those notions often yield almost everywhere solutions (meaning Lipschitz functions satisfying the Hamilton–Jacobi equation almost everywhere). In the case of a Hamiltonian $H(x, p)$ which is convex in the momentum variable p , it was proved by Zhukovskaya (see [Zhu93] and also [BC06, Wei13a]) that both notions provide the same solution. However, they may differ in the general case. Indeed, variational solutions do not necessarily verify the semi–group (or Markov) property. There is a simple way (proposed by Chaperon) to force variational solutions to verify this property and Wei [Wei13a] recently established that the obtained functions are the viscosity solutions (see also the work of Roos [Roo13] on this matter).

In this paper we will focus on the solvability of the following system of uncoupled Hamilton–Jacobi equations:

$$\begin{cases} \partial_t u + H(x, d_x u) = 0 & \text{in } (0, +\infty) \times (0, +\infty) \times M \\ \partial_s u + G(x, d_x u) = 0 & \text{in } (0, +\infty) \times (0, +\infty) \times M \\ u(0, 0, x) = u_0(x) & \text{on } M. \end{cases}$$

Such system is known as *multi–time Hamilton–Jacobi equation* and has its roots in economics. There is now a consequent literature concerning this problem, where either one or the other notion of solution has been considered. The issue about existence and uniqueness of viscosity solutions of the multi–time equation was first addressed by Lions and Rochet [LR86] in the case of convex and coercive Hamiltonians on $\mathbb{R}^n \times \mathbb{R}^n$ only depending on the momentum variable. This work was subsequently generalized in [BT01] to Hamiltonians depending on both variables of $\mathbb{R}^n \times \mathbb{R}^n$, but still convex and coercive in the momentum. As a counterpart, a commutation hypothesis is necessary (which is always verified in the case of Lions and Rochet), namely the vanishing of the Poisson bracket:

$$\{H, G\} := \partial_p H \cdot \partial_x G - \partial_p G \cdot \partial_x H = 0 \quad \text{in } \mathbb{R}^n \times \mathbb{R}^n.$$

Those results were then extended to less regular settings in [MR06].

In the framework of symplectic geometry and variational solutions, Cardin and Viterbo [CV08] extended these results to a larger class of Hamiltonians that do not satisfy any convexity assumption in the momentum variable. The authors proved that the null Poisson bracket condition entails existence and uniqueness of the variational solutions of the multi–time equation.

Knowing the results of Cardin and Viterbo, and the close link between variational and viscosity solutions, it seems natural to ask the following question: if two regular Hamiltonians (non necessarily convex) H and G satisfy the null Poisson bracket condition, can we solve the multi–time Hamilton–Jacobi in the viscosity sense?

The purpose of this note is showing that the answer to this question is, in general, negative.

The article is organized as follows. In the first section we recall the definitions of viscosity and variational solutions and we present the link between the two through Wei’s result. In the second section we give our main result. More precisely, we show

that, when the convexity condition on the Hamiltonians is dropped, the multi-time equation admits a viscosity solution only for very special choices of the initial datum u_0 (see Theorem 2.1 for the precise statement). Given a pair of non-convex Hamiltonians H and G , the corresponding set of admissible initial data is typically empty. An example of this fact is provided in Corollary 2.4.

In the appendix we give the proof of a folklore theorem on existence and uniqueness of viscosity solutions to the evolutionary Hamilton–Jacobi equation. The result is proved under a set of conditions either more general or different than the ones usually assumed in the literature. The techniques and ideas employed are by no means new, however, since they are disseminated in the literature and the adaptation to the case at issue requires some technical work, we felt it might be useful to gather them in a single reference and provide a neat proof of this fact.

1. PRELIMINARIES

Throughout the paper, we will call *Hamiltonian* a continuous function defined on the cotangent bundle of a smooth and compact Riemannian manifold M . We will say that H is *coercive* if

$$\lim_{|p| \rightarrow \infty} H(x, p) = +\infty \quad \text{for every } x \in M. \quad (\text{C})$$

For convenience, we will assume that M is the flat n -dimensional torus \mathbb{T}^n , however all the results keep holding in the more general setting since they are essentially local in nature.

In this section, we focus on the case of a single Hamilton–Jacobi equation, i.e.

$$\partial_t u + H(x, D_x u) = 0 \quad \text{in } (0, +\infty) \times \mathbb{T}^n, \quad (\text{HJ})$$

where we assume either H or $-H$ to be a coercive Hamiltonian. When H is at least C^1 with Lipschitz derivatives ($C^{1,1}$ in short), it induces a vectorfield, called *Hamiltonian vectorfield*, defined as follows:

$$X_H(x, p) = (\partial_p H(x, p), -\partial_x H(x, p)) \quad \text{for all } (x, p) \in \mathbb{T}^n \times \mathbb{R}^n.$$

When well defined, the flow of this vectorfield will be denoted by (φ^t) .

1.1. Viscosity solutions. Let u be a continuous function in $(0, +\infty) \times \mathbb{T}^n$.

The function u is called a *viscosity subsolution* of (HJ) if, for every function $\varphi \in C^1((0, +\infty) \times \mathbb{T}^n)$ such that $u - \varphi$ attains a local maximum at $(t, x) \in (0, +\infty) \times \mathbb{T}^n$, we have

$$\partial_t \varphi(x, t) + H(x, D_x \varphi(x, t)) \leq 0. \quad (1.2)$$

Any such test function φ will be called *supertangent* to u at (t, x) .

The function u is called a *viscosity supersolution* of (HJ) if, for every function $\varphi \in C^1((0, +\infty) \times \mathbb{T}^n)$ such that $u - \varphi$ attains a local minimum at $(t, x) \in (0, +\infty) \times \mathbb{T}^n$, we have

$$\partial_t \varphi(x, t) + H(x, D_x \varphi(x, t)) \geq 0. \quad (1.3)$$

Any such test function φ will be called *subtangent* to u at (t, x) . Last, u is called a *viscosity solution* of (HJ) if it is both a sub and a supersolution.

It is well known, see for instance [Bar94], that the notions of viscosity sub and supersolutions are local, in the sense that the test function φ needs to be defined only in a neighborhood of the point (t, x) . Moreover, up to adding to φ a quadratic term, such a point can be always assumed to be either a strict maximum or a strict

minimum point of $u - \varphi$. In this instance, we will say that φ is a *strict supertangent* (resp., *strict subtangent*) to u at (t, x) .

One of the main features of the notion of viscosity solution is that it is extremely stable. Indeed, in rough terms, if \tilde{u} is a small perturbation of u and the test function φ is, for instance, a strict subtangent to u at (t, x) , then it will also be a subtangent to \tilde{u} at a point (\tilde{t}, \tilde{x}) close to (t, x) . It follows that a continuous function obtained as local uniform limit of a sequence of viscosity solutions of (HJ) is still a solution of the same equation. Furthermore, it has the advantage to be a reasonable notion, in the sense that it yields existence and uniqueness results under mild conditions, as it is illustrated by the following now classical result

Theorem 1.1. *Let $H : \mathbb{T}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that either H or $-H$ is a coercive Hamiltonian. If $u_0 \in C(\mathbb{T}^n)$, there exists a unique uniformly continuous function u in $[0, +\infty) \times \mathbb{R}^n$ satisfying $u(0, \cdot) = u_0$ and solving (HJ) in the viscosity sense. Furthermore, if u_0 is Lipschitz-continuous in \mathbb{T}^n , then u is Lipschitz continuous in $[0, +\infty) \times \mathbb{R}^n$ and its Lipschitz constant only depends on $\|Du_0\|_\infty$ and H .*

The proof of this result can be always reduced to the case when the Hamiltonian is coercive thanks to the following simple remark:

Proposition 1.2. *Let u be a continuous function in $(0, +\infty) \times \mathbb{T}^n$. Then u is a subsolution (respectively, a supersolution) of (HJ) if and only if $-u$ is a supersolution (resp., a subsolution) of*

$$\partial_t v - H(x, -D_x v) = 0 \quad \text{in } (0, +\infty) \times \mathbb{T}^n.$$

Theorem 1.1 is known by PDE specialists, however all the the proofs we have found in literature are given under sets of assumptions somewhat different from the ones herein considered. For completeness, we have provided in the Appendix a proof of Theorem 1.1 in a slightly more general form, namely we deal with the case when \mathbb{T}^n is replaced by \mathbb{R}^n , see Theorem 2.5. Theorem 1.1 follows from this by taking the lift of H and of the initial datum u_0 to the universal cover of the torus.

In the rest of the paper, we will always assume the initial datum u_0 to be Lipschitz in \mathbb{T}^n and we will denote by $S_H^t u_0(x)$, or simply by $S^t u_0(x)$ when no ambiguity is possible, the unique Lipschitz solution $u(t, x)$ to the evolutionary Hamilton–Jacobi equation (HJ) with Hamiltonian H and taking the initial datum u_0 at $t = 0$.

We end this section by recalling some facts about Tonelli Hamiltonians. We remind that a Hamiltonian H is termed *Tonelli* if it is of class C^2 , satisfies the following superlinear growth in p :

$$\frac{H(x, p)}{|p|} \xrightarrow{|p| \rightarrow \infty} +\infty,$$

and is strictly convex, in the sense that $\partial_{pp}^2 H$ is everywhere positive definite as a quadratic form. We will be interested in some regularity properties of solutions to the Hamilton–Jacobi equation. Recall that a function $f : \mathbb{T}^n \rightarrow \mathbb{R}$ is said to be locally semi-concave (resp. semi-convex) (with a linear modulus) if it can be locally written as the sum of a smooth function and a concave (resp. convex) function (see [CS04] for a complete presentation). In particular, a locally semi-concave function has a supertangent at every point. This implies, for example, that the map $x \mapsto |x|$ is not locally semi-concave because of its singularity at 0 which is downward. The following well-known fact holds, (see for example Lemma 4.2 in [Ber12]):

Lemma 1.3. *Let H be a Tonelli Hamiltonian and u_0 an initial datum. Then, for all $t > 0$, the function $S^t u_0$ is locally semi-concave in \mathbb{T}^n .*

We also underline for further use that a function is both locally semi-concave and locally semi-convex if and only if it is $C^{1,1}$ (C^1 with Lipschitz derivative, see Theorem 6.1.5 in [Fat08]).

Using this idea, Patrick Bernard proved the following result (Proposition 4.10 in [Ber12]):

Proposition 1.4. *Let u_0 be a semi-convex function in \mathbb{T}^n and H a Tonelli Hamiltonian. Then, for $t > 0$ small enough, $S^t u_0$ is $C^{1,1}$.*

Finally, we will need the following beautiful result due to Fathi (Theorem 6.4.1 in [Fat08]):

Theorem 1.5. *Let $v : \mathbb{T}^n \rightarrow \mathbb{R}$ be a C^1 function and H be a Tonelli Hamiltonian whose flow will be denoted (φ^t) . Let L be the graph of the differential of v . If there is an increasing and diverging sequence $(t_n)_{n \in \mathbb{N}}$ such that $\varphi^{t_n}(L)$ is a graph above \mathbb{T}^n for all $n \in \mathbb{N}$, then L is invariant by (φ^t) . Moreover, it follows that v is $C^{1,1}$ and that there exists a constant $\alpha[0] \in \mathbb{R}$, only depending on H , such that v is a strong solution of $H(x, D_x v) = \alpha[0]$ in \mathbb{T}^n . In particular*

$$S^t v(x) = v(x) - t\alpha[0] \quad \text{for all } (t, x) \in [0, +\infty) \times \mathbb{T}^n.$$

The constant $\alpha[0]$ appearing above is called *critical value* for H . It is characterized as the only constant a for which an equation of the kind $H(x, D_x v) = a$ in \mathbb{T}^n admits viscosity solutions, see [LPV87]. The viscosity solutions of the corresponding critical equation were used by Fathi as central objects in the weak KAM and Aubry-Mather theory (see [Fat08, Fat12, Ber12] for introductions), hence they are also known as *weak KAM solutions*.

1.2. Variational solutions and their link with viscosity solutions. In this section we give a very brief overview of the notion of variational solutions. For a more detailed account we refer to [BC06, Wei13a] and to the references therein.

We will assume here that $H : \mathbb{T}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is a C^2 Hamiltonian with compact levels, meaning that $H^{-1}(\{a\})$ is compact for all $a \in \mathbb{R}$. This ensures that H has what is called finite propagation speed and that all the notions that will be used are well defined. Moreover, its Hamiltonian flow (φ^t) is complete (indeed, H remains constant along the trajectories of the flow).

Let us consider a smooth function $u_0 : \mathbb{T}^n \rightarrow \mathbb{R}$, and we will denote by $L_{u_0} \subset \mathbb{T}^n \times \mathbb{R}^n$ the graph of the differential of u_0 . It is a smooth Lagrangian submanifold of $\mathbb{T}^n \times \mathbb{R}^n$ for its standard symplectic structure. As it is commonly done, we introduce the Hamiltonian

$$\mathcal{H} : \mathbb{R} \times \mathbb{T}^n \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}, \quad (t, x, \tau, p) \mapsto \tau + H(x, p),$$

so that solving the evolutionary Hamilton–Jacobi equation is equivalent to solving $\mathcal{H}(t, x, D_{(t,x)} U) = 0$. We also define

$$\Gamma_{u_0} = \{(0, x, -H(x, p), p), \quad (x, p) \in L_{u_0}\}.$$

Let us denote by (Φ^t) the Hamiltonian flow generated by $X_{\mathcal{H}} = (\partial_{(\tau,p)} \mathcal{H}, -\partial_{(t,x)} \mathcal{H})$. For every fixed $T > 0$ we set

$$L_{\mathcal{H}, u_0} = \bigcup_{t \in [0, T]} \Phi^t(\Gamma_{u_0}).$$

It is then known that $L_{\mathcal{H},u_0}$ admits a *generating function quadratic at infinity*, that is a function

$$S : [0, T] \times \mathbb{T}^n \times \mathbb{R}^k \rightarrow \mathbb{R}, \quad (t, x, \eta) \mapsto S(t, x, \eta),$$

for some integer k , such that S coincides with a quadratic form outside a compact set, 0 is a regular value of the mapping $(t, x, \eta) \mapsto \partial_\eta S(t, x, \eta)$ and

$$L_{\mathcal{H},u_0} = \left\{ (t, x, \partial_{(t,x)} S(t, x, \eta)) : \partial_\eta S(t, x, \eta) = 0 \right\}.$$

Moreover, the construction of S is done in such a way that if $\partial_\eta S(0, x, \eta) = 0$ then $S(0, x, \eta) = u_0(x)$. Then, define

$$V^t u_0(x) := \inf_{[\sigma]=A} \max_{\eta \in \sigma} S(t, x, \eta),$$

where A generates some homology group depending on the topology of the sublevels of S . The function $(t, x) \mapsto V^t u_0(x)$ is the variational solution. This construction can also be generalized to Lipschitz initial data by limit arguments.

If H is convex in p , then $S^t u_0(x) = V^t u_0(x)$ for any Lipschitz continuous initial datum u_0 . However, for general Hamiltonians, this is not always the case since $V^t \circ V^s u_0$ may differ from $V^{t+s} u_0$. This is what is commonly referred to as the lack of the *semi-group property*. Recently, Wei [Wei13a, Wei13b] established the following link (which is formulated in a more general way in her thesis):

Theorem 1.6 ([Wei13a]). *For any Lipschitz function $u_0 : \mathbb{T}^n \rightarrow \mathbb{R}$, the following holds:*

$$S^t u_0(x) = \lim_{n \rightarrow +\infty} (V^{t/n})^n u_0(x) \quad \text{for all } (t, x) \in (0, +\infty) \times \mathbb{T}^n.$$

2. THE MULTI-TIME HAMILTON-JACOBI EQUATION

We will only be interested in systems of two equations. Therefore, in the following, H and G will be two smooth and proper Hamiltonians on $\mathbb{T}^n \times \mathbb{R}^n$. Note that no convexity assumptions are taken in the momentum variable. Given a Lipschitz function $u_0 : \mathbb{T}^n \rightarrow \mathbb{R}$, we are looking for a real function $u(t, s, x)$ defined on $[0, +\infty) \times [0, +\infty) \times \mathbb{T}^n$ which solves

$$\begin{cases} \partial_t u + H(x, D_x u) = 0 & \text{in } (0, +\infty) \times (0, +\infty) \times \mathbb{T}^n \\ \partial_s u + G(x, D_x u) = 0 & \text{in } (0, +\infty) \times (0, +\infty) \times \mathbb{T}^n \\ u(0, 0, x) = u(x) & \text{on } \mathbb{T}^n \end{cases} \quad (2.1)$$

in the viscosity sense. As first remarked in [BT01], a necessary condition for viscosity solutions of (2.1) to exist, for any Lipschitz initial datum, is that H and G commute, in the sense that their Poisson bracket vanishes, i.e.

$$\{H, G\} := \partial_p H \cdot \partial_x G - \partial_p G \cdot \partial_x H = 0 \quad \text{in } \mathbb{T}^n \times \mathbb{R}^n.$$

This can be derived by differentiating the equation and by using the method of characteristics (for smooth initial data), see for instance Appendix C in [DZ13]. When the Hamiltonians are convex and coercive in the momentum variable, this condition is sufficient as well, as it is proved in [BT01]. In the more general case when the Hamiltonians are proper but not necessarily convex in the momentum, it is proved in [CV08] that the null Poisson bracket condition implies the existence of variational solutions to the multi-time equation. Knowing the links between variational and viscosity solutions (Theorem 1.6), it seems natural to wonder if the

vanishing of $\{H, G\}$ also implies the existence of solutions to (2.1) in the viscosity sense. Using the semi-group property and Theorem 1.1, it is easily seen that this is equivalent to requiring that, for any Lipschitz $u_0 : \mathbb{T}^n \rightarrow \mathbb{R}$, the following hold:

$$u(t, s, \cdot) = S_H^t \circ S_G^s u_0 = S_G^s \circ S_H^t u_0 \quad \text{for every } s, t > 0.$$

We will prove the following theorem:

Theorem 2.1. *Let H and G be two commuting Hamiltonians on $\mathbb{T}^n \times \mathbb{R}^n$ such that both H and $-G$ are Tonelli. Then there exists a solution to the multi-time equation (2.1) with Lipschitz initial datum $u_0 : \mathbb{T}^n \rightarrow \mathbb{R}$ if and only if u_0 is $C^{1,1}$ and the graph of its differential is invariant by both the Hamiltonian flows of H and G . Moreover, in this case, there are two constants c_H and c_G such that*

$$H(x, D_x u_0) = c_H \quad \text{and} \quad G(x, D_x u_0) = c_G, \quad \text{for every } x \in \mathbb{T}^n,$$

which means that

$$S_H^t \circ S_G^s u_0 = S_G^s \circ S_H^t u_0 = u_0 - t c_H - s c_G \quad \text{for every } s, t > 0.$$

Remark 2.2. Note that $c_H = \alpha_H[0]$ and $c_G = -\alpha_{-G}[0]$, where $\alpha_H[0]$ and $\alpha_{-G}[0]$ are the critical values associated with H and $-G$.

Before proving this theorem, we establish a simple lemma:

Lemma 2.3. *Let $G : \mathbb{T}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that either G or $-G$ is a coercive Hamiltonian. Let us set*

$$\overline{G}(x, p) := -G(x, -p) \quad \text{for every } (x, p) \in \mathbb{T}^n \times \mathbb{R}^n.$$

Then, for any Lipschitz function $u_0 : \mathbb{T}^n \rightarrow \mathbb{R}$, the following holds :

$$S_G^s u_0 = -S_{\overline{G}}^s(-u_0) \quad \text{for every } s > 0.$$

In particular, if $-G$ is Tonelli, then $S_G^s u$ is locally semi-concave for any $s > 0$.

Proof. The first assertion follows from Proposition 1.2, while the second is a consequence of Lemma 1.3. \square

In the next proof, recall that if $w : \mathbb{T}^n \rightarrow \mathbb{R}$ is any C^1 function, we denote by $L_w \subset \mathbb{T}^n \times \mathbb{R}^n$ the graph of its differential.

Proof of Theorem 2.1. Let u_0 be an initial condition for which there exists a solution to the multi-time equation. We have seen that in this case

$$S_G^s \circ S_H^t u_0 = S_H^t \circ S_G^s u_0 \quad \text{for every } s, t > 0.$$

In particular, as the left hand side is locally semi-concave by Lemma 2.3 and the right hand side is locally semi-convex, by Lemma 1.3, we deduce that for all $s, t > 0$, the function $S_G^s \circ S_H^t u_0 = S_H^t \circ S_G^s u_0$ is $C^{1,1}$.

Let $s > 0$ be fixed. For $\varepsilon > 0$, set $v_\varepsilon = S_G^s \circ S_H^\varepsilon u_0 \in C^{1,1}$. It follows from [Fat08, Proposition 4.11.1 and Theorem 6.2.2] that

$$\varphi_H^t(L_{v_\varepsilon}) = L_{S_H^t v_\varepsilon} \quad \text{for every } t > 0.$$

In particular, we may let t go to infinity and use Fathi's theorem 1.5, yielding the existence of c_H such that

$$S_H^t v_\varepsilon = v_\varepsilon - t c_H \quad \text{for every } t \geq 0.$$

Now letting $\varepsilon \rightarrow 0^+$ and passing to the limit, we deduce that

$$S_H^t \circ S_G^s u_0 = S_G^s u_0 - t c_H \quad \text{for every } t \geq 0. \quad (2.2)$$

By continuity, the above identity holds for $s = 0$ as well.

Let us now fix $t > 0$. The same argument applied to \overline{H} , \overline{G} and $-u_0$ yields the existence of a constance c_G such that

$$S_G^s \circ S_H^t u_0 = S_H^t u_0 - s c_G \quad \text{for every } s \geq 0. \quad (2.3)$$

By continuity, the above identity holds for $t = 0$ as well. Putting (2.2) and (2.3) together we get

$$S_H^t \circ S_G^s u_0 = u_0 - s c_G - t c_H = S_G^s \circ S_H^t u_0 \quad \text{for every } s, t \geq 0,$$

finally showing that u_0 is $C^{1,1}$ and a strong solution of the stationary equations associated with H and with G . \square

Even if for a Tonelli Hamiltonian H there are always weak KAM solutions, they may all fail to be $C^{1,1}$. In this instance, the previous theorem implies that there are no global solutions to the multi-time equation for H and $-H$ (nor in fact for H and $-G$ for any Tonelli Hamiltonian G).

We furnish an example to conclude.

Corollary 2.4. *There exists a Tonelli Hamiltonian $H : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$ such that the multi-time equation (2.1) with $G := -H$ does not admit viscosity solutions, for any Lipschitz initial datum u_0 .*

Proof. According to Theorem 2.1, we just need to exhibit a Tonelli Hamiltonian H that does not admit weak KAM solutions of class $C^{1,1}$. The example that follows is classical, see for instance [Fat08, Section 4.14]. Consider the pendulum equation, corresponding to

$$H(x, p) = \frac{1}{2}|p|^2 + \cos(2\pi x), \quad \text{on } \mathbb{T}^1 \times \mathbb{R}.$$

The Hamiltonian vector-field is then given by $X_H(x, p) = (p, 2\pi \sin(2\pi x))$. It possesses 2 fixed points, $(\frac{1}{2}, 0)$ which is an elliptic fixed point, and $(0, 0)$ which is hyperbolic. The stable and unstable manifold are given by the two separatrix which equations are $p = \pm \sqrt{2(1 - \cos(2\pi x))} = \pm 2 \sin(\pi x)$. Those separatrix bound an elliptic island filled with periodic trajectories. In this case, it is known that $\alpha[0] = 1$. Moreover, the only weak KAM solution (up to addition of a constant) is given by

$$u(x) = \begin{cases} \int_0^x 2 \sin(\pi t) dt = \frac{2}{\pi}(1 - \cos(\pi x)), & \forall x \in [0, \frac{1}{2}], \\ \frac{2}{\pi} - \int_{\frac{1}{2}}^x 2 \sin(\pi t) dt = \frac{2}{\pi}(1 + \cos(\pi x)), & \forall x \in (\frac{1}{2}, 1), \end{cases}$$

which is not of class $C^{1,1}$. Indeed, the graph of the derivative of any other weak KAM solution v must stay on the union of the two separatrix, the fact that v is semi-concave means that this graph may only "jump downwards", therefore it may possess only one discontinuity, and in order for v to be 1-periodic, this jump must occur at $1/2$. Hence $v' = u'$, that is $v - u$ is constant. \square

It is proved in [Zav10, CL11] that two commuting Tonelli Hamiltonians have the same weak KAM solutions, therefore the previous theorem is consistent with this fact, in the sense that it is normal that whenever H has a $C^{1,1}$ weak KAM solution, it will be the same for G .

Let us conclude with one last speculation. The previous results, in order to be global, are stated for Tonelli Hamiltonians. However, when a Hamiltonian is convex

in some areas, and concave in others, the previous discussions show that according to where the characteristics come from, the variational solutions of the evolutionary equation are either semi-concave or semi-convex in some zones. Those two different kinds of regularization might give a new insight as to why variational solutions differ from viscosity solutions, and do not verify the semi-group property.

APPENDIX

In this Appendix, we want to give a proof of Theorem 1.1 in a slightly more general setting. In what follows we will denote by $UC(X)$ and $BUC(X)$ the space of uniformly continuous and bounded uniformly continuous real functions on the metric space X , respectively.

We consider the following evolutionary Hamilton–Jacobi equation:

corretto tempo

$$\partial_t u + F(x, D_x u) = 0 \quad \text{in } (0, +\infty) \times \mathbb{R}^n, \quad (\text{A})$$

where the Hamiltonian $F : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the following conditions:

- (F1) $F \in BUC(\mathbb{R}^n \times B_R)$ for every $R > 0$;
- (F2) $\inf_{x \in \mathbb{R}^n} F(x, p) \rightarrow +\infty$ as $|p| \rightarrow +\infty$.

We will prove the following result:

Theorem 2.5. *Let $F : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ be a Hamiltonian satisfying (F1)-(F2) and let $u_0 \in UC(\mathbb{R}^n)$. Then there exists a unique function $u \in UC([0, +\infty) \times \mathbb{R}^n)$ with $u(0, \cdot) = u_0$ that solves (A) in the viscosity sense. If u_0 is Lipschitz in \mathbb{R}^n , then u is Lipschitz continuous in $[0, +\infty) \times \mathbb{R}^n$ and its Lipschitz constant only depends on $\|Du_0\|_\infty$ and F .*

We need to generalize the notions of viscosity sub and supersolution given at the beginning of Section 1.1 to the case of discontinuous functions. Let Ω be an open subset of $(0, +\infty) \times \mathbb{R}^n$ and let $u : \Omega \rightarrow \mathbb{R}$ be a possibly discontinuous locally bounded function. We define the *upper* and *lower semicontinuous envelope* of u , denoted by u^* and u_* respectively, as follows:

$$u^*(t, x) := \limsup_{(s, y) \rightarrow (t, x)} u(s, y), \quad u_*(t, x) := \liminf_{(s, y) \rightarrow (t, x)} u(s, y)$$

for every $(t, x) \in \Omega$. We say that u^* is a (*non-continuous*) *viscosity subsolution* of (A) in Ω if (1.2) holds for any supertangent φ to u^* at the point $(t, x) \in \Omega$. The gradient of any such supertangent φ at the point (t, x) is called *supergradient* of u^* at (t, x) . We will denote by $D^+u^*(t, x)$ the set consisting of all supergradients of u^* at (t, x) . Analogously, we say that u_* is a *supersolution* of (A) in Ω if (1.3) holds for any subtangent φ to u_* at the point $(t, x) \in \Omega$.

We will need the following comparison result between discontinuous sub and supersolutions.

Proposition 2.6. *Let v and u be, respectively, an upper and lower semicontinuous function on $[0, +\infty) \times \mathbb{R}^n$. Let us assume that, for every $\alpha > 0$, there exists a constant $M_\alpha > 0$ such that, for every $(t, x) \in (0, +\infty) \times \mathbb{R}^n$,*

- (v_0) $v(t, x) \geq v(0, x) - (\alpha + M_\alpha t)$,
- (u_0) $u(t, x) \geq u(0, x) - (\alpha + M_\alpha t)$.

If v is a subsolution and u is a supersolution of (A), we have

$$\sup_{[0,+\infty) \times \mathbb{R}^n} (v - u) \leq \sup_{\mathbb{R}^n} (v(0, \cdot) - u(0, \cdot)).$$

Remark 2.7. While condition (u_0) is necessary for the comparison principle, see for instance Example 4.9 in [CS05], the analogous condition (v_0) is redundant. It is assumed to avoid further technical details. We remark that conditions (u_0) and (v_0) are always fulfilled if u and v are both absolutely continuous in $[0, +\infty) \times \mathbb{R}^n$. In particular, Proposition 2.6 entails uniqueness of the solution of (A) in $\text{UC}([0, +\infty) \times \mathbb{R}^n)$.

For the proof of Proposition 2.6, we will need to regularize the subsolution v through the following *sup-convolution in time*, see [BCD97]: for every fixed $\delta > 0$, define

$$v^\delta(t, x) := \sup_{s \geq 0} \left\{ v(s, x) - \frac{|t - s|}{\delta} \right\}, \quad (t, x) \in [0, +\infty) \times \mathbb{R}^n. \quad (2.4)$$

For every $(t, x) \in [0, +\infty) \times \mathbb{R}^n$, we denote by $\mathcal{M}^\delta(t, x)$ the set of $s \geq 0$ that realize the supremum in the definition of $v^\delta(t, x)$. We set

$$\Omega^\delta := \text{int} \left\{ (t, x) \in (0, +\infty) \times \mathbb{R}^n : v^\delta(t, x) > v(0, x) - \frac{t}{\delta}, \right\},$$

that is, Ω^δ is the largest open subset of $(0, +\infty) \times \mathbb{R}^n$ made up of points (t, x) for which $0 \notin \mathcal{M}^\delta(t, x)$. Last, we set $\mu := \max\{1, \sup_{\mathbb{R}^n \times \mathbb{R}^n} -F\}$, which is finite by assumption (F2). The following holds:

Proposition 2.8. *Let us assume that v is upper semicontinuous on $[0, +\infty) \times \mathbb{R}^n$ and a non-continuous subsolution of (A). Then for every $\delta < 1/\mu$ we have:*

- (i) $v(t, x) \leq v^\delta(t, x) \leq v(0, x) + \mu t$ for every $(t, x) \in [0, +\infty) \times \mathbb{R}^n$.
- (ii) For every $(t, x) \in [0, +\infty) \times \mathbb{R}^n$ the set $\mathcal{M}^\delta(t, x)$ is nonempty and contained in $[0, 2t/(1 - \delta\mu)]$. In particular, v^δ is upper semicontinuous in $[0, +\infty) \times \mathbb{R}^n$.
- (iii) If $(t_0, x_0) \in \Omega^\delta$, then

$$D^+v^\delta(t_0, x_0) \subseteq D^+v(s_0, x_0) \quad \text{for every } s_0 \in \mathcal{M}^\delta(t_0, x_0).$$

In particular, v^δ is an upper semicontinuous subsolution of (A) in Ω^δ .

- (iv) The function $v^\delta(\cdot, x)$ is $1/\delta$ -Lipschitz continuous in $[0, +\infty)$ for every fixed $x \in \mathbb{R}^n$. Moreover, there exists a positive constant L_δ such that

$$|D_x v^\delta(t, x)| \leq L_\delta \quad \text{for a.e. } (t, x) \in \Omega^\delta.$$

- (v) Let us assume that v satisfies (v) for some positive constants α and M_α . Then for every $\delta > 0$ small enough

$$\left[2\alpha \frac{\delta}{1 - \delta M_\alpha}, +\infty \right) \times \mathbb{R}^n \subset \Omega^\delta$$

Proof. (i) The first inequality is obvious. To prove the second inequality, we first observe that v satisfies

$$\partial_t v \leq -F(x, Dv) \leq \mu \quad \text{in } (0, +\infty) \times \mathbb{R}^n$$

in the viscosity sense. We infer

$$v(s, x) \leq v(\tau, x) + \mu(s - \tau) \quad \text{for every } x \in \mathbb{R}^n \text{ and } 0 < \tau < s,$$

in particular, by letting $\tau \rightarrow 0^+$ and by using the upper semicontinuity of v , we get

$$v(s, x) \leq v(0, x) + \mu s \quad \text{for every } (s, x) \in [0, +\infty) \times \mathbb{R}^n. \quad (2.5)$$

For $\delta < 1/\mu$ we get

$$v^\delta(t, x) \leq v(0, x) + \sup_{s \geq 0} \left\{ \mu s - \frac{|t-s|}{\delta} \right\} = v(0, x) + \mu t.$$

(ii) Let us fix $(t, x) \in [0, +\infty) \times \mathbb{R}^n$. The fact that $\mathcal{M}^\delta(t, x)$ is nonempty follows from (2.5) and the upper semicontinuity of v . Pick $s \in \mathcal{M}^\delta(t, x)$. From the definition of $v^\delta(t, x)$ and (2.5) we have

$$v(0, x) - \frac{t}{\delta} \leq v^\delta(t, x) = v(s, x) - \frac{|s-t|}{\delta} \leq v(0, x) + \mu s - \frac{|s-t|}{\delta}.$$

A straightforward computation shows that $s \leq 2t/(1 - \delta\mu)$. The upper semicontinuity of v^δ readily follows from this and from the upper semicontinuity of v .

(iii) First note that $(t_0, x_0) \in \Omega^\delta$ implies $\mathcal{M}^\delta(t_0, x_0) \subset (0, +\infty)$. To prove the first assertion, it suffices to observe that, if φ is a supertangent to v^δ at $(t_0, x_0) \in \Omega^\delta$, then the function $(t, x) \mapsto \varphi(t + (t_0 - s_0), x)$ is a supertangent to v at (s_0, x_0) for every fixed $s_0 \in \mathcal{M}^\delta(t_0, x_0)$. Since v^δ is upper semicontinuous by (ii) and F is time-independent, we infer that v^δ is an upper semicontinuous subsolution of (A) in Ω^δ .

(iv) The fact v^δ is $1/\delta$ -Lipschitz in t is apparent by definition. By using (iii) we get

$$F(x, Dv^\delta) \leq -\partial_t v^\delta \leq \frac{1}{\delta} \quad \text{in } \Omega^\delta,$$

and the estimate on $|D_x v^\delta|$ in Ω^δ follows from the coercivity of F , see Lemma 2.5 in [Bar94].

(v) Pick $\delta < 1/M_\alpha$ and set $T_\delta := 2\delta\alpha/(1 - \delta M_\alpha)$. Then every $(t, x) \in (T_\delta/2, +\infty) \times \mathbb{R}^n$ satisfies

$$v(0, x) - \frac{t}{\delta} < v(0, x) - (M_\alpha t + \alpha) \leq v(t, x) \leq v^\delta(t, x),$$

in particular $[T_\delta, +\infty) \times \mathbb{R}^n \subset \Omega^\delta$, as it was to be shown. \square

Proof of Proposition 2.6. The proof is divided in two steps.

Step 1. We first assume v Lipschitz continuous in $[0, +\infty) \times \mathbb{R}^n$, i.e. there exists $L > 0$ such that

$$|v(t, x) - v(s, y)| \leq L(|x - y| + |t - s|) \quad \text{for every } (t, x), (s, y) \in [0, +\infty) \times \mathbb{R}^n.$$

We define a function $\Phi : ([0, +\infty) \times \mathbb{R}^n)^2 \rightarrow \mathbb{R}$ by setting

$$\Phi(t, x, s, y) := v(t, x) - u(s, y) - \frac{1}{2\varepsilon} (|x - y|^2 + |t - s|^2) - \beta(\langle x \rangle + \langle y \rangle) - \eta(t^2 + s^2),$$

where $\langle x \rangle := \sqrt{1 + |x|^2}$, η is a positive constant that will be suitably chosen, and ε, β are positive parameters that will be sent to 0.

We set $A := \sup_{\mathbb{R}^n} (v(0, \cdot) - u(0, \cdot)) < +\infty$ and we assume that $A < +\infty$. Let us suppose by contradiction that there exists $(\hat{t}, \hat{x}) \in (0, +\infty) \times \mathbb{R}^n$ such that

$$v(\hat{t}, \hat{x}) - u(\hat{t}, \hat{x}) = A + \gamma \quad \text{for some } \gamma > 0.$$

We choose $\eta > 0$ and $\beta > 0$ small enough such that

$$2\beta\langle\hat{x}\rangle + 2\eta\hat{t}^2 \leq \frac{\gamma}{2}.$$

Then

$$\sup_{([0,+\infty)\times\mathbb{R}^n)^2} \Phi \geq \Phi(\hat{t}, \hat{x}, \hat{t}, \hat{x}) = A + \gamma - (2\beta\langle\hat{x}\rangle + 2\eta\hat{t}^2) \geq A + \frac{\gamma}{2}$$

We proceed to show that such a supremum is attained. Indeed, choose $0 < \alpha < \gamma/4$ and $M = M(\alpha) > 0$ such that

$$u(s, y) \geq u(0, y) - (\alpha + Ms) \quad \text{for every } (s, y) \in [0, +\infty) \times \mathbb{R}^n.$$

Then

$$\begin{aligned} \Phi(t, x, s, y) &\leq A + L(t + |x - y|) + (\alpha + Ms) \\ &\quad - \frac{1}{2\varepsilon} (|x - y|^2 + |t - s|^2) - \beta(\langle x \rangle + \langle y \rangle) - \eta(t^2 + s^2). \end{aligned} \quad (2.6)$$

It is easily seen that $\Phi(t, x, s, y) \rightarrow -\infty$ when $(|x| + |y|) + (t + s) \rightarrow +\infty$, so there exists $(\bar{t}_\varepsilon, \bar{x}_\varepsilon, \bar{s}_\varepsilon, \bar{y}_\varepsilon) \in ([0, +\infty) \times \mathbb{R}^n)^2$ such that

$$\Phi(\bar{t}_\varepsilon, \bar{x}_\varepsilon, \bar{s}_\varepsilon, \bar{y}_\varepsilon) = \max_{([0,+\infty)\times\mathbb{R}^n)^2} \Phi \geq A + \frac{\gamma}{2} \quad (2.7)$$

This maximum point also depends on the parameters η and β , but such dependence is omitted to ease notations.

Claim 1: there exists $\varepsilon_0 > 0$ such that

$$\bar{t}_\varepsilon + \bar{s}_\varepsilon \geq \frac{\gamma}{4(L + M)} =: \theta \quad \text{for every } \varepsilon \leq \varepsilon_0.$$

We make use of (2.7) and exploit the estimate (2.6). Since for all positive Y , the following holds

$$Y \left(L - \frac{Y}{2\varepsilon} \right) \leq L^2 \frac{\varepsilon}{2},$$

we get (with $Y = |\bar{x}_\varepsilon - \bar{y}_\varepsilon|$)

$$A + \frac{\gamma}{2} \leq \Phi(\bar{t}_\varepsilon, \bar{x}_\varepsilon, \bar{s}_\varepsilon, \bar{y}_\varepsilon) \leq A + (L + M)(\bar{t}_\varepsilon + \bar{s}_\varepsilon) + \alpha + L^2 \frac{\varepsilon}{2}$$

i.e.

$$(L + M)(\bar{t}_\varepsilon + \bar{s}_\varepsilon) \geq \frac{\gamma}{2} - \left(\alpha + L^2 \frac{\varepsilon}{2} \right).$$

Since $\alpha < \gamma/4$, the claim follows provided $\varepsilon_0 > 0$ is chosen small enough.

Claim 2: $\frac{|\bar{t}_\varepsilon - \bar{s}_\varepsilon|^2}{\varepsilon} \leq \frac{L^2 + M^2}{\eta}$ for every $\varepsilon \leq \varepsilon_0$.

From (2.7) and (2.6) and by arguing similarly as above we get

$$\begin{aligned} A + \frac{\gamma}{2} &\leq A + \alpha + \bar{t}_\varepsilon(L - \eta\bar{t}_\varepsilon) + \bar{s}_\varepsilon(M - \eta\bar{s}_\varepsilon) + \frac{\varepsilon}{2}L^2 - \frac{|\bar{t}_\varepsilon - \bar{s}_\varepsilon|^2}{2\varepsilon} \\ &\leq A + \alpha + \frac{L^2}{4\eta} + \frac{M^2}{4\eta} + \frac{\varepsilon}{2}L^2 - \frac{|\bar{t}_\varepsilon - \bar{s}_\varepsilon|^2}{2\varepsilon}, \end{aligned}$$

i.e.

$$\frac{|\bar{t}_\varepsilon - \bar{s}_\varepsilon|^2}{2\varepsilon} \leq \frac{L^2}{4\eta} + \frac{M^2}{4\eta} - \left(\frac{\gamma}{2} - \left(\alpha + \frac{\varepsilon}{2}L^2 \right) \right) < \frac{L^2 + M^2}{2\eta}$$

This shows that $|\bar{t}_\varepsilon - \bar{s}_\varepsilon| \rightarrow 0$ as $\varepsilon \rightarrow 0^+$, hence, by Claim 1, $\bar{t}_\varepsilon > 0$, $\bar{s}_\varepsilon > 0$ for $\varepsilon > 0$ small enough. We can now exploit the fact that $\varphi(t, x) := \Phi(t, x, \bar{s}_\varepsilon, \bar{y}_\varepsilon)$ is

a supertangent to v at $(\bar{t}_\varepsilon, \bar{x}_\varepsilon) \in (0, +\infty) \times \mathbb{R}^n$ and $\psi(s, y) := \Phi(\bar{t}_\varepsilon, \bar{x}_\varepsilon, s, y)$ is a sub-
 tangente to u at $(\bar{s}_\varepsilon, \bar{y}_\varepsilon) \in (0, +\infty) \times \mathbb{R}^n$. Since v is an upper semicontinuous
 subsolution and u is a lower semicontinuous supersolution of (A), we get

$$\partial_t \varphi(\bar{t}_\varepsilon, \bar{x}_\varepsilon) - \partial_s \psi(\bar{s}_\varepsilon, \bar{y}_\varepsilon) \leq F(\bar{y}_\varepsilon, D\psi(\bar{s}_\varepsilon, \bar{y}_\varepsilon)) - F(\bar{x}_\varepsilon, D\varphi(\bar{t}_\varepsilon, \bar{x}_\varepsilon)) \quad (2.8)$$

An easy computation shows that

$$\partial_t \varphi(\bar{t}_\varepsilon, \bar{x}_\varepsilon) - \partial_s \psi(\bar{s}_\varepsilon, \bar{y}_\varepsilon) = 2\eta(\bar{t}_\varepsilon + \bar{s}_\varepsilon) \geq 2\eta\theta,$$

where the inequality follows from Claim 1, while

$$|D\varphi(\bar{t}_\varepsilon, \bar{x}_\varepsilon) - D\psi(\bar{s}_\varepsilon, \bar{y}_\varepsilon)| \leq 2\beta.$$

Moreover, by the fact that v is L -Lipschitz and φ is a supertangent to v at $(\bar{t}_\varepsilon, \bar{x}_\varepsilon)$, we get

$$|D\varphi(\bar{t}_\varepsilon, \bar{x}_\varepsilon)| = \left| \frac{\bar{x}_\varepsilon - \bar{y}_\varepsilon}{\varepsilon} + \beta \frac{\bar{x}_\varepsilon}{\langle \bar{x}_\varepsilon \rangle} \right| \leq L,$$

yielding in particular that $|\bar{x}_\varepsilon - \bar{y}_\varepsilon| \rightarrow 0$ as $\varepsilon \rightarrow 0^+$. We now send ε and β to 0 in (2.8) and we make use of (F1) and of the above inequalities to get $2\eta\theta \leq 0$, in contrast with the definition of θ .

Step 2. Let us prove the result in the general case. Let us fix $\alpha > 0$ and let us accordingly choose $M_\alpha > 0$ in such a way that condition (v_0) holds. Let us set $T_\delta := 2\delta\alpha/(1 - \delta M_\alpha)$. According to Proposition 2.8, for $\delta > 0$ small enough the function $(t, x) \mapsto v^\delta(T_\delta + t, x)$ is a Lipschitz continuous subsolution of (A) in $[0, +\infty) \times \mathbb{R}^n$, hence we can apply Step 1 to infer

$$\sup_{[0, +\infty) \times \mathbb{R}^n} (v^\delta(T_\delta + \cdot, \cdot) - u) \leq \sup_{\mathbb{R}^n} (v^\delta(T_\delta, \cdot) - u(0, \cdot)).$$

Moreover, we know that

$$v(t, x) - \frac{T_\delta}{\delta} \leq v^\delta(t, x) - \frac{T_\delta}{\delta} \leq v^\delta(T_\delta + t, x) \leq v(0, x) + \mu(T_\delta + t),$$

so

$$\sup_{[0, +\infty) \times \mathbb{R}^n} (v - u) \leq \sup_{\mathbb{R}^n} (v(0, \cdot) - u(0, \cdot)) + \frac{2\alpha}{1 - \delta M_\alpha} (1 + \mu\delta).$$

The assertion follows by sending $\delta \rightarrow 0^+$ and since $\alpha > 0$ was arbitrarily chosen. \square

The proof of Theorem 2.5 is standard. It follows from Proposition 2.6 by making use of Perron's method.

Proof of Theorem 2.5. We only need to show the existence of a solution, for the uniqueness comes from Proposition 2.6, see also Remark 2.7.

Let us first assume $u_0 \in \text{Lip}(\mathbb{R}^n)$. In view of the assumptions (F1)-(F2), there exists a positive constant M such that $M > \sup\{|F(x, p)| : |p| \leq \|Du_0\|_\infty\}$. Then the functions

$$\underline{u}(t, x) := u_0(x) - Mt, \quad \bar{u}(t, x) := u_0(x) + Mt, \quad (t, x) \in [0, +\infty) \times \mathbb{R}^n$$

are, respectively, a sub and a supersolution to (A). We denote by \mathcal{S} the family of continuous functions on $[0, +\infty) \times \mathbb{R}^n$ that are subsolutions of (A) and we set, for every $(t, x) \in [0, +\infty) \times \mathbb{R}^n$,

$$u(t, x) = \sup \{v(t, x) : v \in \mathcal{S}, \underline{u} \leq v \leq \bar{u} \text{ on } [0, +\infty) \times \mathbb{R}^n\}.$$

Then u^* and u_* are, respectively, a sub and a supersolution of (A), see Theorem 2.14 in [BCD97], and satisfy $u^*(0, \cdot) = u_*(0, \cdot) = u_0$ on \mathbb{R}^n . By applying the comparison

principle stated in Proposition 2.6, we infer $u^* \leq u_*$ on $[0, +\infty) \times \mathbb{R}^n$. Since the other inequality is obvious by definition, we get that u is continuous on $[0, +\infty) \times \mathbb{R}^n$, solves (A) in the viscosity sense and takes the initial datum u_0 at $t = 0$. It is left to show the Lipschitz continuity of u . By the comparison principle and the definition of u we get

$$\|u(t+h, \cdot) - u(t, \cdot)\|_{L^\infty(\mathbb{R}^n)} \leq \|u(h, \cdot) - u(0, \cdot)\|_{L^\infty(\mathbb{R}^n)} \leq Mh.$$

In other words, the function u is M -Lipschitz in t . Using the fact that u is a subsolution of (A), we get that u satisfies

$$F(x, D_x u) \leq -\partial_t u \leq M \quad \text{in } (0, +\infty) \times \mathbb{R}^n$$

in the viscosity sense. By the coercivity of F , we conclude that $u(t, \cdot)$ is Lipschitz for every $t > 0$, with a constant only depending on u_0 and F , see for instance Lemma 2.5 in [Bar94].

If $u_0 \in \text{UC}(\mathbb{R}^n)$, we can find, for instance by sup-convolution, see [BCD97], a sequence of Lipschitz functions $u_0^k : \mathbb{R}^n \rightarrow \mathbb{R}$ that uniformly converge to u_0 on \mathbb{R}^n . Let us denote by $u^k : [0, +\infty) \times \mathbb{R}^n \rightarrow \mathbb{R}$ the corresponding Lipschitz solution of (A) with initial datum u_0^k . By the comparison principle we have

$$\|u^m - u^k\|_{L^\infty([0, +\infty) \times \mathbb{R}^n)} \leq \|u_0^m - u_0^k\|_{L^\infty(\mathbb{R}^n)},$$

that is, $(u^k)_k$ is a Cauchy sequence in $[0, +\infty) \times \mathbb{R}^n$ with respect to the sup-norm. Hence the Lipschitz functions u_k uniformly converge to a function u on $[0, +\infty) \times \mathbb{R}^n$, which is therefore uniformly continuous. By the stability of the notion of viscosity solution, we conclude that u is a solution of (A) with initial datum u_0 . \square

ACKNOWLEDGMENT

The second author wishes to thank Patrick Bernard for inciting him to write this note and Valentine Roos for explaining him the content and details of her work.

REFERENCES

- [Bar94] G. Barles. *Solutions de viscosité des équations de Hamilton-Jacobi*, volume 17 of *Mathématiques & Applications (Berlin) [Mathematics & Applications]*. Springer-Verlag, Paris, 1994.
- [BC06] O. Bernardi and F. Cardin. Minimax and viscosity solutions of Hamilton-Jacobi equations in the convex case. *Commun. Pure Appl. Anal.*, 5(4):793–812, 2006.
- [BCD97] M. Bardi and I. Capuzzo-Dolcetta. *Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations*. Systems & Control: Foundations & Applications. Birkhäuser Boston Inc., Boston, MA, 1997. With appendices by Maurizio Falcone and Pierpaolo Soravia.
- [Ber12] P. Bernard. The Lax-Oleinik semi-group: a Hamiltonian point of view. *Proc. Roy. Soc. Edinburgh Sect. A*, 142(6):1131–1177, 2012.
- [BT01] G. Barles and A. Tourin. Commutation properties of semigroups for first-order Hamilton-Jacobi equations and application to multi-time equations. *Indiana Univ. Math. J.*, 50(4):1523–1544, 2001.
- [Cha91] M. Chaperon. Lois de conservation et géométrie symplectique. *C. R. Acad. Sci. Paris Sér. I Math.*, 312(4):345–348, 1991.
- [CL83] M. G. Crandall and P.-L. Lions. Viscosity solutions of Hamilton-Jacobi equations. *Trans. Amer. Math. Soc.*, 277(1):1–42, 1983.
- [CL11] X. Cui and J. Li. On commuting Tonelli Hamiltonians: autonomous case. *J. Differential Equations*, 250(11):4104–4123, 2011.

- [CS04] P. Cannarsa and C. Sinestrari. *Semiconcave functions, Hamilton-Jacobi equations, and optimal control*. Progress in Nonlinear Differential Equations and their Applications, 58. Birkhäuser Boston Inc., Boston, MA, 2004.
- [CS05] F. Camilli and A. Siconolfi. Time-dependent measurable Hamilton-Jacobi equations. *Comm. Partial Differential Equations*, 30(4-6):813–847, 2005.
- [CV08] F. Cardin and C. Viterbo. Commuting Hamiltonians and Hamilton-Jacobi multi-time equations. *Duke Math. J.*, 144(2):235–284, 2008.
- [DZ13] A. Davini and M. Zavidovique. Weak KAM theory for nonregular commuting Hamiltonians. *Discrete Contin. Dyn. Syst. Ser. B*, 18(1):57–94, 2013.
- [Fat08] A. Fathi. Weak KAM Theorem in Lagrangian Dynamics (preliminary version n. 10), Lyon, 15 June 2008.
- [Fat12] A. Fathi. Weak KAM from a PDE point of view: viscosity solutions of the Hamilton-Jacobi equation and Aubry set. *Proc. Roy. Soc. Edinburgh Sect. A*, 142(6):1193–1236, 2012.
- [LPV87] P.-L. Lions, G. Papanicolaou, and S.R.S. Varadhan. Homogenization of Hamilton-Jacobi equation. unpublished preprint, 1987.
- [LR86] P.-L. Lions and J.-C. Rochet. Hopf formula and multitime Hamilton-Jacobi equations. *Proc. Amer. Math. Soc.*, 96(1):79–84, 1986.
- [MR06] M. Motta and F. Rampazzo. Nonsmooth multi-time Hamilton-Jacobi systems. *Indiana Univ. Math. J.*, 55(5):1573–1614, 2006.
- [Roo13] V. Roos. Solutions de l'équation de Hamilton-Jacobi. Thèse de Master 2, ENS, 2013.
- [Wei13a] Q. Wei. Solutions de viscosité des équations de Hamilton-Jacobi et minmax itérés. Thèse de Doctorat, Université de Paris 7, 2013.
- [Wei13b] Q. Wei. Viscosity solution of Hamilton-Jacobi equation by a limiting minmax method. *ArXiv e-prints*, March 2013.
- [Zav10] M. Zavidovique. Weak KAM for commuting Hamiltonians. *Nonlinearity*, 23(4):793–808, 2010.
- [Zhu93] T. Zhukovskaya. Singularités de minimax et solutions faibles d'équations aux dérivées partielles, 1993.

DIP. DI MATEMATICA, SAPIENZA UNIVERSITÀ DI ROMA, P.LE ALDO MORO 2, 00185 ROMA, ITALY

E-mail address: `davini@mat.uniroma1.it`

IMJ (PROJET ANALYSE ALGÈBRIQUE), UPMC, 4, PLACE JUSSIEU, CASE 247, 75252 PARIS CÉDEX 5, FRANCE

E-mail address: `zavidovique@math.jussieu.fr`